

ON THE RELATIVE STABILITY OF CUMULENONE AND ALDEHYDE ISOMERS: WHEN WE HEAT345(Q) THINGS UP

KELVIN LEE, *Radio and Geoastronomy Division, Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA*; MICHAEL C MCCARTHY, *Atomic and Molecular Physics, Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA*; JOHN F. STANTON, *Department of Chemistry, The University of Texas, Austin, TX, USA*.

Isomers of $\text{H}_2\text{C}_{2n+1}\text{O}$ are examples of complex organic molecules that are either known or proposed to exist in the interstellar medium. For the smallest of these chains ($\text{H}_2\text{C}_3\text{O}$) only two of three isomers are observed in space: propynal ($\text{HC}(\text{O})\text{CCH}$) and cyclopropenone ($\text{c} - \text{C}_3\text{H}_2\text{O}$), while evidence for the remaining isomer propadienone ($\text{H}_2\text{C}_3\text{O}$) is currently lacking. Potentially, this behaviour may be rationalised by a thermodynamic argument: several studies have provided quantum chemical calculations in an effort to determine the relative thermodynamic stability between these three isomers. An early study by Radom, at the SCF/6-31G** level ranked $\text{HC}(\text{O})\text{CCH}$ as the thermodynamic minimum, followed by $\text{H}_2\text{C}_3\text{O}$, and $\text{c} - \text{C}_3\text{H}_2\text{O}$. The most recent determination by Karton and Talbi, using W2-F12 theory, places $\text{H}_2\text{C}_3\text{O}$ as the lowest energy isomer; 2.5 kJ mol^{-1} lower than the $\text{HC}(\text{O})\text{CCH}$ form. In an attempt to resolve this long-standing ambiguity, we were motivated to provide high level calculations based on the HEAT protocol. In this talk, we will discuss the relative stability of $\text{H}_2\text{C}_3\text{O}$ and $\text{H}_2\text{C}_5\text{O}$ isomers, along with their sulfur analogues, as revealed by HEAT345(Q) theory.